References

[1] Ali. A., S. M. Al-Ogaily, N. A. Al-Asgah and J. Gropp: 2003. Effect of sub-lethal concentrations of copper on the growth performance of *Oreochromis niloticus. J. Appl. Ichthyol.*, 19: 183-188. (para no. 15)

[2] Allen. P: 1995. Accumulation profiles of cadmium and their modification with mercury and lead in the edible tissues of Oreochromis aureus. *Fresh Environ Bull* 2:745-751, (para no.1)

[3] Amundsen. P. A., F. J. Staldvik, A. A. Lukin, N. A. Kashulin, O.A. Popova and Y. S. Reshetnikov: 1997. Heavy metal contamination in fresh water fish from the border region between Norway and Russia. *Sci. Total Environ*., 201: 111-124. (para no. 17)

[4] Ashraj. W: 2005. Accumulation of heavy metals in kidney and heart tissues of Epinephelus microdon fish from the Arabian Gulf. Environmental Monitoring Assessment, 1-3(103), pp 311-316. (para no. 3)

[5] Ashraj. W: 2005. Accumulation of heavy metals in kidney and heart tissues of *Epinephelus microdon* fish from the Arabian Gulf. Environ. Monit. Assess. 101(1-3): 311-316. (para no. 10)

[6] Ayfer. Y. and L. Jacobs: 1995. Synergestic effects of temperature, oxygen and water flow on accumulation and tissue distribution of mercury in carp (*Cyprinus carpio* L.). *Chemosphere*, 31 (11-12), 4437-4453 (para no. 16)

[7] Azmat. H. M. Javed and G. Jabeen: 2012. Acute toxicity of aluminium to the fish *(Catla catla, Labeo rohita and Cirrhina mrigala). Pakistan Vet. J.*, 32: 85-87. (para no. 14)

[8] Basa, Siraj. P, Usha Rani. A: 2003. Cadmium induced antioxidant defense mechanism in freshwater teleost *Orechromis mossambicus* (tilapia). EcoToxicology Environment Safety, 2(56), pp 218-221. (para no. 3)

[9] Biswas. K., Chattopadhyay. I., Banerjee. R. K. and Bandyopadhyay. U: 2002. Biological activities and medicinal properties of neem (*Azadirachta indica*). Current Science, 82: 1336-1334 (para no. 5)

[10] Buhl. J: 1997. Relatives sensitivity of three endangered fishes, Colorado squa fish, bony tail, and razor back sucker, to selected metal pollutants. Ecotoxicol. Environ. Saf, 37(2), 186-192 (para no. 16)

[11] Canli. M. A y, O. Kalay. M: 1998. Levels of heavy metals (Cd, Pb, Cu and Ni) in tissue of *Cyprinus carpio, Barbus capito* and *Chondrostoma regium* from the Seyhan river. Turkish Journal of Zoology, 3(22), pp 149-157. (para no. 3)

[12] Chattopadhyay S, Anam K, Aditya A K: 1995. Time dependent distribution of heavy metal in different tissues of Labeo rohita *J Freshwat Biol* 7: 203-205, (para no.1)

[13] David. M, Mushier S. B and Philip G H: 2003. Alterations in the levels in tissues of fresh water fish Labeo rohitha exposed to Fenvalerate Polin.; 22(3): 359-369. (para no. 11)

[14] Dirilgen. N: 2001. Accumulation of heavy metals in fresh water organisms: Assessment of toxic interactions.FAO.Fischer.Technology, 212, pp 1-13. (para no. 3)

[15] Eisler. R: 1998. Copper Hazards to Fish. Wildlife and Invertebrates: A Synoptic Review. Biological Science Repot I 'SGSBRD'BRS-1997-0002. (para no.8)

[16] Farkas. A., Salanki J. and Specziar A: 2002. Relation between growth and heavy metal concentration in organs of bream, *Abramis brama* L, population lake Balaton. Arch. Environ. Contam. Toxicol., 43: 236-243. (para no.7)

[17] Farombi. E. O, Adelowo O. A., Ajimoko Y R: 2007. Biomarkers of oxidatives stress and heavy metal levels as indicators of environmental pollution in African Cat Fish (*Clarias gariepinus)* from Nigeria Ogun River, Int. J. Environ. Res. Pub. Health, 4(2): 158-165. (para no. 10)

[18] Farombi. E. O, O.A. Adelowo and R. K. Ajimoko: 2007. Biomarkers of oxidative stress and heavy metal levels as indicators of environmental pollution in African Cat Fish

(*Clarias gariepinus)* from Nigeria Ogun River, *Int. J. Environ. Res. Pub. Health*, 4: 158-165. (para no. 17)

[19] FSY: 2010. Fisheries Statistical Yearbook of Bangladesh, 24th edition. Department of Fisheries, Ministry of Fisheries and livestock, government of the People's Republic of Bangladesh. (para no.18)

[20] Goyer. R. A: 1991. Toxic effects of metals In: Amer, M.O., Doul J Klaassen C. D. (Ed) Casarett and Doull's Toxicology; Basic science of poisons, Pergmon Press, Oxford. (para no.7)

[21] Gupta. P. and N. Srivastava: 2006. Effects of sub-lethal concentration of zinc on histological changes and bioaccumulation of zinc by kidney of fish *Channa punctatus* (Bloch). *J. Environ. Biol.,* 27: 211-215. (para no.15)

[22] Gupta. A. K. and V. K. Rajbanshi: 1991. Toxicity of copper and cadmium to *Heteropneustes fossilis* (Bloch*). Acta Hydrochim. Et. Hydroboil.,* 19(3), 331-340 (para no. 16)

[23] Hussain. M. G. & Mazid. M. A: 2005. Carp Genetic Resources of Bangladesh In : Penman D. J., Gupta M. V. and Dey M. M. (eds.), Carp Genetic Resources for Aquaculture in Asia, World Fish Center Technical Report, 65, 152pp. (para no.18)

[24] ICAR: 1993. World Neem Conference Souvenir ICAR, Banglore, India (para no.5)

[25] Isani. G., G. Andreani, F. Cocchioni, D. Fedeli, E. Carpene and G. Falcioni: 2009. Cadmium accumulation and biochemical responses in Sparus aurata following sub lethal Cd exposure. *Ecotoxical. Environ. Saf.* 72: 224-230. (para no.17)

[26] Javed. M and S Abudullah: 2006. Studies on the acute and lethal toxicities of iron and nickel to fish. *Pakistan J. Biol. Sci*. 9: 330-335. (para no.17)

[27] John. P. R., V. Rena and D J. McQueen: 1996. Uptake rates of food-chain and waterborne mercury by fish: Field measurement, a mechanistic model and an assessment of uncertainties. *Can. J. Fish. Aquat. Sci.,* 53(2), 395-407 (para no. 16)

[28] Joseph. T. A., S, M. Sebastin and S. Raja: 1992. Toxicity of nickel on protein content in tissue of *Cyprinus carpio* (Linn). Indian J. Environ. Hlth, 34(3), 236-238 (para no. 13)

[29] Kalay. M. and M. Canli: 2000. Elimination of essential (Cu, Zn) and non-essential (Cd, Pb) metals from tissues of freshwater fish *Tilapia zill.* Turkish. J. Zool., 24: 429-436. (para no. 14)

[30] Khare. S and S. Singh: 2002. Histopathological lesions induced by copper sulphate and lead nitrate in the gills of fresh fish Nandus. J. *Ecotoxicol. Environ. Monit.,* 12: 105-111. (para no.17)

[31] Kock G., Triendl M. & R. Hofer: 1996. Seasonal patterns of metal accumulation in Artic char (Salvelinus alpinus) from an oligotrophic Alpine lake related to temperature. Can. J. Fish. Aquat. Sci. 53: 780-786 (para no.7)

[32] Kumar Lca, Vincent S, Ambrose T: 1994. Uptake and persistence of the heavy metal cadmium in tissues of the fresh water fish *Cyprinus carpio. Poll Res* 13: 361-364, (para no.1)

[33] Leonard. A. W., R. V, Hyne, R. Lim, F. Pablo and P. Van den Brink: 2000. Riverline endosulfan concentrations in the Namoi River, Australia: Link to cotton field runoff and microinvertebrate population densities. Enviro. Toxical. Chem., 19, 1540-1551 (para no.9)

[34] Loh. K. C., T. S. Chung and A. Wei-Fern: 2000. Immobilized cell membrane bioreactor for high strength phenol waste water. *J. Environ. Eng.,* 126, 75-79 (para no. 12)

[35] Mathieson. S., S. G George and D. S. Mclusky: 1996. Temporal variations of total mercury concentrations and burdens in the liver of eel pout Zoarces viviparous from the fourth estuary, Scotland: Implications for mercury biomonitoring. *Mar. Ecol. Prog. Ser.,* 138, 41-49 (para no. 16)

[36] McCrary. J. E. and M. G. Heagler: 1997. The use of simultaneous multiple species acute toxicity tests to compare the relative sensitivities of aquatic organisms to mercury. *J. Environ. Sci. health (Part A). Environ. Sci. Eng. & Toxic & Hazard. Subst. Cont.*, 32(1), 73-81 (para no. 16)

[37] Miah. M. I., Harun. M. S., Rahman. M. M: 2009. The Embryonic and Larvel Development of an Endangered Species of Bata *(Labeo bata). Int. J. Sustain. Corp Prod.*, 4(1): 72-82. (para no.20)

[38] Naeem. M., A. Salam, S. S. Tahir and N. Rauf: 2011. The effect of fish size and condition on the contents of twelve essential and non essential elements in *Aristichthys nobilis. Pakistan Vet. J.* 31: 109-112. (para no.15)

[39] Naz. S., M. Javed, S. Hayat, S. Abdullah, M. Bilal and T. Shaukat: 2008. Long term effect of lead (Pb) toxicity on the growth performance, nitrogen conversion ratio and yields of major carps. *Pakistan J. Agric. Sci.* 45: 53-58. (para no.15)

[40] Olaifa, F. E, Olaifa A. K, Adelaja. A. A, Owolabi. A. G: 2004, Heavy metal contamination of *Clarias gariepinus* from a Lake and Fish in Ibadan, Nigeria. African Journal of Biomedical Research. 7, pp 145-148. (para no. 3)

[41] Patel J M., Bahadur A: 2011. Histopathological Manifestations of Sub Lethal toxicity of Copper Ions in *Catla catla*. Am. Eur. J. Toxicol. Sci. 3(1): 01-05. (para no. 10)

[42] Rajan. M. R: 1990. Sub lethal effects of textile mill effluent on protein, carbohydrate and lipid content of different tissues of fish *Cyprinus carpio. Environ. Ecol.*, 8(1), 54-58 (para no. 13)

[43] Ramaneshwari. K. and L. M. Rao: 2008. Influence of endosulfan and monocrotophos exposure in the activity of NADPH cytochrome C reductase (NCCR) of *Labeo rohita* (Ham). J. Environ. Bio, 29, 183-185 (para no.9)

[44] Samir A M, Zaahkouk Eman G E, Hetal Talaat E I, Abd-Rabo and Somania Z A Rashed: 2000.Carbamate toxicity and protective effect of vit-A and vit-E: Somebio-

chemical aspects of male albino rats. The Egyptian J Hospital Medicine. 1: 60-77. (para no. 11)

[45] Sarkar. U. K. and Larka, W. S: 2010. Small indigenous freshwater fish species of India: Significance, conservation and utilization. *Aquaculture Asia Magazine* Vol. XV (3): 34-35.

[46] Sastry K V, Shukla V: 1994. Influence of protective agents in the toxicity of cadmium to a fresh water fish (*Channa punctatus*) *Bull Environ Contam Toxicol* 53: 711-717, (para no.1)

[47] Saxsena R, Garg P: 2011. Vitamin E provides protection against *In vitro* oxidative stress due to pesticide (Chlorphrifos and Endosulfan) in goat RBC, GERF Bull. Biosci. Article, press. (para no. 10)

[48] Srividya. K and Mohanty. K: 2009. Biosorption of hexavalent chromium from aqueous solution by *Catla catla* scales: Equilibrium and kinetics studies. *Chemical Engineering Journal,* 155: 666-673. (para no.19)

[49] Subramanian. M. A. and G. Varadaraj: 1993. Impact of tannery effluent on the biochemical constituents in the haemolymph of the freshwater prawn. *Macrobrachium idella* Heller. *J. Environ. Biol.,* 14(3), 255-259 (para no. 13)

[50] Sukhovaskaya. I. V., L. P. Smimov, N. N. Nemova and V. T. Komov: 2001. Effect of mercurium on the fractional composition of the low-molecular peptides of musculature in the perch, *Perca flaviatillis. Voprosy Ikhtiol*., 41(5), 699-703 (para no. 16)

[51] Velez. D., Montoro. R: 1998. Arsenic speciation in manufactured seafood products: a review. Journal of good Protect, 9(61), pp 1240-1245. (para no. 3)

[52] Vinodhini R, Muthuswamy N: 2008. Effect of heavy metals induced toxicity on metabolic biomarkers in common carp. *Cyprinus carpio.* Maejo Int. J. Sci. Technol. 2: 192-200. (para no. 10)

[53] Vosyliene. M, Z. Jankaite, A: 2006. Effect of heavy metal model mixture on rainbow trout biological parameters Ekologija, 4, pp 12-17. (para no. 3)

[54] Vosyliene. M, Z. A. Jankaite: 2006. Effect of heavy metal model mixture on rainbow trout biological parameters *Ekologija*, Nr. 4: 12-17. (para no. 15)

[55] Vuren. V. J. H. J., H. H. D. Preez. V. Wepener, A. Anendrofe and I. E.J.K. Barnhoorn: 1999. Lethal and sub-lethal effects of metals on the physiology of fish. An experimental approach with monitoring support. Water research commission, Pretoria. WRC. Report 608/11/99. (para no. 14)

[56] Vutukuru. S. S: 2005. Acute effects of Hexavalent chromium on survival, oxygen consumption, hematological parameters and some biochemical profiles of the Indian major carp, Labeo rohita, International Journal of Environmental Research and Public Health, 2(3), pp 456-462. (para no. 3)

[57] Vutukuru. S. S: 2005. Acute effects of Hexavalent chromium on survival, oxygen consumption, hematological parameters and some biochemical profiles of the Indian Major carp, L. rohita, Int. J. Environ. Res. and Public Health, 2(3), 456-462. (para no. 10)

[58] Waqar. A: 2006. Levels of selected heavy metals in Tuna fish. Arab Journal of Science and Engineering, I A (31), pp 89-92. (para no. 3)